

PATIENT NAME: _____

PROCEDURE DATE: _____ at _____ with _____

Please arrive to the Toronto Immune and Digestive Health Institute at: _____
700 Lawrence Ave. W., 3rd floor, Suite 360, Toronto, ON, M6A 3B4 (T: 647-812-2113)
Enter through the East Tower Entrance, located between Booster Juice and the Foot Institute

Arrange for a pick-up **2 hours after your procedure start time.**

If you do not have an adult accompany you home, your procedure will be cancelled.

The Colonoscopy Package includes the following documents:*

1. Colonoscopy Preparation Instructions (2 pages)
2. Pre-Anesthesia Questionnaire (3 pages)
3. Colonoscopy & Polypectomy Information (2 pages)

*Contact the office immediately if you are missing any of the documents listed above.

Please read **ALL** the documents enclosed within the Colonoscopy Package carefully. Failure to follow the instructions may result in the cancellation of your procedure.

If you have any questions or require further information, please contact the clinic and one of our staff members will be happy to assist you.

Toronto Immune and Digestive Health Institute
700 Lawrence Avenue West
3rd floor, Suite 360,
Toronto, ON M6A 3B4
Phone: 647-812-2113
Email: reception@tidhi.ca

CANCELLATION POLICY

You will be called by our office two weeks prior to your scheduled procedure to remind you of your appointment date and time. If our office is not able to speak with you directly, you **must** call the clinic at 647-812-2113 no later than 5 business days prior to your scheduled procedure. Failure to confirm your appointment in a timely manner may result in your procedure time being given away to another patient.

CANCELLATIONS WITH LESS THAN 5 BUSINESS DAY'S NOTICE WILL RESULT IN A \$200 FEE NOT COVERED BY OHIP

PLEASE READ THE FOLLOWING INSTRUCTIONS CAREFULLY

MEDICATIONS TO DISCONTINUE:

- **Iron medications** – stop at least **10** days prior to the procedure
- **Aspirin** – stop at least **7** days prior to your procedure, after consulting with your physician
- **Coumadin (Warfarin) or Xarelto (Rivaroxaban)** – you may need to temporarily discontinue these medications. Please discuss this with your prescribing physician well in advance of the procedure
- **Insulin or pills for Diabetes** – please discuss with your family physician prior to the test
- If you have an abnormal heart valve, prosthetics heart valve or other conditions where you have been told to take antibiotics prior to dental procedures, please advise us immediately.

1. TWO WEEKS PRIOR to your procedure:

- Call the clinic at 647-812-2113 to confirm your procedure
- Complete the Pre-Anesthesia Questionnaire and submit to the clinic via email (reception@tidhi.ca) or fax (**647-812-2114**)
- Purchase the following bowel prep materials over-the-counter at your local pharmacy:
 - (1) box of **Pico-Salax** (contains two sachets)
 - (1) box of **Dulcolax 5mg** tablets

2. ONE WEEK PRIOR to your procedure:

- STOP eating whole grains, nuts, seeds, dried fruit, or raw fruits/vegetables

3. ONE DAY PRIOR to your procedure:

- You must be on **clear fluids ALL DAY**
 - DO NOT DRINK OR EAT any solid or milk dairy products
 - DO NOT DRINK any fluids with red or purple colour.
 - **Clear fluids include:** soft drinks (sprite, 7-UP, Ginger Ale), clear broth, fruit juices (without pulp), water, tea or coffee (without milk or cream), jell-o, ice popsicles
- **At 5:00pm:** Take **two Dulcolax tablets** with water
- **At 8:00pm:** Take **first dose of Pico-Salax**
 - Empty the contents of 1 sachet of Pico-Salax into a cup. Add 150 mL of cold water and stir frequently for 2 -3 minutes. Sometimes the reaction of mixing PICO-SALAX and cold water will cause the solution to become hot. It is becomes hot, wait until it cools sufficiently to drink.
 - Drink an additional 2 litre (as tolerated) of clear fluids after finishing the first sachet

4. DAY OF PROCEDURE:

- **At 6:00am:** Take **second dose** of **Pico-Salax** (same instructions as first dose)
 - Drink plenty of clear fluids (as tolerated) of clear fluids after finishing the second sachet
- Take all of your usual medications (except those listed on page 1) with small sips of water.
- **DO NOT EAT ANYTHING ON THE DAY OF THE PROCEDURE**
- **DO NOT DRINK ANYTHING FOUR HOURS PRIOR TO THE PROCEDURE**

IMPORTANT INFORMATION

- If you have sleep apnea plan to bring your CPAP machine to the procedure
- Continue any blood pressure medication as usual
- If you missed any of the steps on the previous page, **PLEASE STILL COME IN FOR THE PROCEDURE**
- **DO NOT BRING ANY VALUABLES JEWELLERY OR WEAR MAKE UP**
- You will be sedated during the procedure. Therefore, **YOU MUST BE ACCOMPANIED BY SOMEONE WHEN YOU LEAVE.** If you do have someone to accompany you, your procedure will be cancelled.

WHEN YOU ARRIVE

- The Toronto Immune and Digestive Health Institute is located in the Lawrence-Allen Centre at (700 Lawrence Avenue West). There is free, unlimited parking and adjacent TTC/Subway access (Lawrence West Station).
- Please enter through the East Tower entrance, which is accessible from the parking lot on the south side of the mall facing Lawrence Avenue. The East Tower entrance can be found between Booster Juice and the Foot Institute.
- Once inside, take the elevators up to the 3rd floor – Suite 360.
- Unfortunately due to COVID restrictions, your escort cannot accompany you inside the clinic. Your escort can visit the retail shops at the Lawrence-Allen Centre, which provides immediate access to food, shopping and other amenities. Your escort will be informed when you are ready to be picked up.

Please complete this form and email (reception@tidhi.ca) or fax (647-812-2114) a copy of the completed form to the clinic at least 14 days prior to your scheduled procedure.

Date of Procedure (MM/DD/YY):

Name (Last, First, M.I.):		DOB (MM/DD/YY):	
OHIP #:	Version Code:	Weight:	Height:
Emergency Contact Name:	Relationship:	Phone:	

MEDICAL HISTORY

Cardiac Health		
Angina	☐ Yes	☐ No
Hypertension	☐ Yes	☐ No
Myocardial Infarction	☐ Yes	☐ No
Atrial Fibrillation	☐ Yes	☐ No
Heart murmur	☐ Yes	☐ No
Cardiac Valve Disorders	☐ Yes	☐ No
Pacemaker	☐ Yes	☐ No
Stent/Angioplasty	☐ Yes	☐ No
Other:		

Blood Health		
Anemia	☐ Yes	☐ No
Sickle Cell Trait	☐ Yes	☐ No
Sickle Cell Anemia	☐ Yes	☐ No
Bleeding Disorder	☐ Yes	☐ No
DVT or PE	☐ Yes	☐ No
Aneurysm	☐ Yes	☐ No
HIV/AIDS	☐ Yes	☐ No
Other:		

Respiratory Health		
Asthma	☐ Yes	☐ No
COPD	☐ Yes	☐ No
Tuberculosis	☐ Yes	☐ No
Sleep Apnea ☐ CPAP	☐ Yes	☐ No
Other:		

Endocrine and Metabolic Health			
Diabetes ☐ Type I ☐ Type II	☐ Yes	☐ No	
Thyroid Disease name:	☐ Yes	☐ No	
Other:			

Gastrointestinal Health		
Heartburn or GERD	☐ Yes	☐ No
Hiatus Hernia	☐ Yes	☐ No
Liver Disease name:	☐ Yes	☐ No
Inflammatory Bowel Disease	☐ Yes	☐ No
Difficulty Eating or Swallowing	☐ Yes	☐ No
Nausea or Vomiting	☐ Yes	☐ No
Other:		

Neuro and Musculoskeletal Health		
Dementia	☐ Yes	☐ No
Alzheimer's disease	☐ Yes	☐ No
Migraine	☐ Yes	☐ No
Vertigo	☐ Yes	☐ No
Neuropathy	☐ Yes	☐ No
Fibromyalgia	☐ Yes	☐ No
Spinal Stenosis	☐ Yes	☐ No
Osteoarthritis	☐ Yes	☐ No
Rheumatoid Arthritis	☐ Yes	☐ No
Epilepsy or Seizure Disorder	☐ Yes	☐ No
Multiple Sclerosis	☐ Yes	☐ No
Parkinson's disease	☐ Yes	☐ No
Other:		

Kidney and Bladder Health		
Kidney Disease <i>name:</i>	☐ Yes	☐ No
If yes, are you on dialysis?	☐ Yes	☐ No
Other:		

Other Significant Conditions:

CURRENT MEDICATIONS

☐ No prescription medications	☐ No over-the-counter medications, supplements, vitamins or probiotics
--	---

Medications, supplements, vitamins or probiotics:	Dose:	Frequency:	Indication:

ALLERGIES

Medication or Substance (<i>e.g. latex, food, etc.</i>):	Type of Reaction:

ANESTHETIC HISTORY

Have you ever had general anesthesia?	☐ Yes	☐ No
<i>If yes, please describe:</i>		
Have you ever had regional anesthesia? (e.g. nerve block, epidural, spinal, etc.)	☐ Yes	☐ No
<i>If yes, please describe:</i>		
Have you had any reactions to anesthesia in the past?	☐ Yes	☐ No <i>If yes, check/describe below.</i>
☐ Malignant Hyperthermia	☐ Pseudocholinesterase	☐ Confusion after surgery
☐ Other Reaction:		
Has a family member ever had a serious reaction to anesthesia?	☐ Yes	☐ No
<i>If yes, please describe:</i>		

ADDITIONAL SCREENING QUESTIONS

Do you bruise easily?	☐ Yes	☐ No
Do you use home oxygen?	☐ Yes	☐ No
Do you have loose teeth or dentures?	☐ Yes	☐ No
Do you have any severe visual or hearing impairments?	☐ Yes	☐ No
Do you drink alcohol?	☐ Yes	☐ No <i>If yes, number of drinks/week:</i>
Do you smoke?	☐ Yes	☐ No <i>If yes, type + amount/week:</i>
Do you use recreational drugs?	☐ Yes	☐ No <i>If yes, type + amount/week:</i>
If female, are you currently pregnant?	☐ Yes	☐ No ☐ N/A

Patient Signature

Date (MM/DD/YY):

BEFORE YOUR PROCEDURE

What is a Colonoscopy?

A colonoscopy is an examination of the large intestine and part of the small intestine with a camera on the end of a tube. Instruments may be passed through the colonoscope to take tiny, painless biopsies (samples of tissue) for microscopic examination in the laboratory. Photographs and video may also be taken during the procedure. Colonoscopy is used to in inflammatory bowel disease (IBD/Crohn's disease, ulcerative colitis) to assess the extent and severity of the inflammation and/or to evaluate treatment response. It is also used to detect and diagnose colon cancer, polyps, causes of bleeding and diarrhea. There is very little risk associated with colonoscopy and polyp removal/biopsies when they are performed by specially trained physicians and health care teams.

What is a Polypectomy?

During the course of your colonoscopy, a polyp may be found. Polyps are abnormal growths of tissue which vary in size from a few millimeters to several centimeters. Depending on the size, shape and location of the polyp, it may be removed (polypectomy). You should feel no pain during polyp removal. Polyps are usually removed because they can be a source of rectal bleeding or may contain cancer. Although the majority of polyps are benign (not cancerous), some may eventually become cancerous if left untouched.

What to Expect

- You will be admitted to the endoscopy suite
- A blood test may be taken
- You will change into a patient gown
- You will empty your bladder
- Your vital signs will be taken
- You will be given an intravenous injection to make you sleepy and relaxed
- The examination will be done with you lying on your left side
- The colonoscope will be introduced via your rectum or stoma so your bowel can be examined
- You may feel some cramping and pressure

AFTER YOUR PROCEDURE

After the Colonoscopy

You will be taken to the Endoscopy recover area where you will remain until most of the effects of the medication have worn off. When your nurse and anesthetist has decided that you are sufficiently recovered, your escort will be called to pick you up. You will be discharged to your escort. You will not be allowed to drive for 24 hours.

Complications

Complications are extremely rare but can occur. One complication is perforation or tear through the wall of the bowel that may allow leakage of intestinal fluids. This complication may require surgery or management with antibiotics and intravenous fluids. Bleeding may occur from a biopsy or polyp removal. This is usually minimal but, rarely, blood transfusions or surgery may be required. Irritation of a vein can occur at an intravenous site. A tender lump or bruise may develop and last from several days to months. It eventually goes away with no ill effects. Other risks include drug reactions and complications from unrelated medical conditions. Your physician and his staff are alert to the possibilities and are fully equipped to handle any complications which may arise.

Results

The results will be explained at the conclusion of your procedure. Because of the sedation, you may not remember what you have been told. If you require further explanation, you should call the physician who performed the procedure or make an office appointment for a complete explanation. If biopsies have been taken, results can take 10 -14 days to be available. Please call the office to follow up.